

References

- Abrams, P.A., 1993. Why predation rate should not be proportional to predator density. *Ecology*, 74: 726 – 733.
- Abrams, P.A., 1994. The fallacies of “ratio-dependent” predation. *Ecology*, 75: 1842 – 1850.
- Abrams, P.A. and L.R. Ginzburg, 2000. The nature of predation: prey dependent, ratio dependent or neither? *Trends in Ecology and Evolution*, 15: 337 – 341.
- Abreu, P.C., C. Hartmann and C. Odebrecht, 1995. Nutrient-rich saltwater and its influence on the phytoplankton of the Patos Lagoon estuary, southern Brazil. *Estuarine, Coastal and Shelf Science*, 40: 219 – 229.
- Adams, S.M., 1976. The ecology of eelgrass, *Zostera marina* (L.), fish communities I: Structural analysis. *Journal of Experimental Marine Biology and Ecology*, 22: 269 – 291.
- Albrecht, K.-F., 1992. Problems of modelling and forecasting on the basis of phenomenological investigations. *Ecological Modelling*, 63: 45 – 69.
- Alongi, D.M., 1998. *Coastal Ecosystem Processes*. CRC Press: New York, London.
- Andersen, K.P. and E. Ursin, 1977. A Multispecies Extension to the Beverton and Holt Theory of Fishing, with Accounts of Phosphorus Circulation and Primary Production. *Meddelelser Fra Danmarks Fiskeri – Og Havundersøgelser*. N.S., 7: 319 – 435.
- Anon, 1996. *Catch and Effort Information Bulletin 1995*. Conservation and Natural Resources: Victorian Fisheries Research Institute: Queenscliff, Victoria, Australia.
- Anon, 1997. *Bays and Inlets Scalefish Fisheries Review: Review of Scalefish Fishing Practices in Victoria’s Bays and Inlets*. WBM Oceanics Australia, Final Report.

- Ansari, Z.A., A. Chatterji, B.S. Ingole, R.A. Sreepada, C.U. Rivonkar and A.H. Parulekar, 1995. Community structure and seasonal variation of an inshore demersal fish community at Goa, west coast of India. *Estuarine, Coastal and Shelf Science*, 41: 593 – 610.
- Arditi, R. and L.R. Ginzburg, 1989. Coupling in predator-prey dynamics: ratio-dependence. *Journal of Theoretical Biology*, 139: 311 - 326
- Arias, A.M. and P. Drake, 1994. Structure and production of the benthic macroinvertebrate community in a shallow lagoon in the Bay of Cádiz. *Marine Ecology Progress Series*, 115: 151 – 167.
- Ariasgonzalez, J.E., O. Hertel and R. Galzin, 1998. Trophic pattern of a reef ecosystem in French Polynesia. *Cybiu*, 22: 159 – 182.
- Bachelet, G., X. de Montaudouin and J-C Dauvin, 1996. The quantitative distribution of subtidal macrozoobenthic assemblages in Arcachon Bay in relation to environmental factors: a multivariate analysis. *Estuarine, Coastal and Shelf Science*, 42: 371 – 391.
- Bardach, J.E., 1959. The summer standing crop of fish on a shallow Bermuda reef. *Limnology and Oceanography*, 4: 77 – 85.
- Baretta, J.W., J.G. Baretta-Bekker and P. Ruardij, 1996. From EMS to ERSEM, towards generic coastal ecosystem models. *Japanese Journal of Aquabiology* 104 (8-3): 197 – 209.
- Baretta, J.W., J.G. Baretta-Bekker and P. Ruardij, 1998. Data needs for ecosystem modelling. *ICES Journal of Marine Science*, 55(4): 756 – 766.
- Baretta, J.W., W. Ebenhöh and P. Ruardij, 1995. The European Regional Seas Ecosystem Model, a complex marine ecosystem model. *Netherlands Journal of*

- Sea Research, 33: 233 – 246.
- Baretta, J.W., P. Ruardij, H.J. Vested and J.G. Baretta-Bekker, 1994. Eutrophication modelling of the North Sea: two different approaches. *Ecological Modelling* 75/76: 471 – 483.
- Baretta-Bekker, J.G. and J.W. Baretta (eds.), 1997. Special Issue: European Regional Seas Ecosystem Model II. *Journal of Sea Research*, 38(3/4).
- Baretta-Bekker, J.G., J.W. Baretta and W. Ebenhöf, 1997. Microbial dynamics in the marine ecosystem ERSEM II with decoupled carbon assimilation and nutrient uptake. *Journal of Sea Research*, 38(3/4): 195 – 211.
- Baretta-Bekker, J.G., J.W. Baretta and K. Rasmussen, 1995. The microbial food web in the European regional Seas ecosystem model. *Netherlands Journal of Sea Research*, 33(3/4): 363 – 379.
- Barnes, R.D., 1987. *Invertebrate Zoology*: 5th ed. Saunders HBJ: Orlando.
- Bartell, S.M., W.G. Cale, R.V. O'Neill and R.H. Gardner, 1988. Aggregation error: research objectives and relevant model structure. *Ecological Modelling*, 41: 157 – 168.
- Barthel, K-G. and R. Goñi, 1995. Preface. *Netherlands Journal of Sea Research*, 33: 229.
- Bax, N. and J.-E. Eliassen, 1990. Multispecies analysis in Balsfjord, northern Norway: solution and sensitivity analysis of a simple ecosystem model. *Journal du Conseil*, 47: 175 – 204.
- Beardall, J., and B. Light, 1997. Microphytobenthos in Port Phillip Bay: Distribution and Primary Productivity. Technical Report No. 30. Port Phillip Bay Environmental Study. CSIRO: Canberra, Australia.
- Beardall, J., S. Roberts and R. Royle, 1996. Phytoplankton in Port Phillip Bay: Spatial

- and seasonal trends in biomass and primary productivity. Technical Report No. 35. Port Phillip Bay Environmental Study. CSIRO: Canberra, Australia.
- Beattie, G., A. Redden and R. Royle, 1996. Microzooplankton Grazing on Phytoplankton in Port Phillip Bay. Technical Report No. 31. Port Phillip Bay Environmental Study. CSIRO: Canberra, Australia.
- Beck, M.B., 1999. Coping with ever larger problems, models, and databases. *Water Science and Technology*, 39: 1 – 11.
- Begon, M. and M. Mortimer, 1986. Population ecology: A unified study of animals and plants. Blackwell Science Publications: London, Melbourne, Boston.
- Bell, J.D., D.A. Pollard, J.J. Burchmore, B.C. Pease and M.J. Middleton, 1984. Structure of a fish community in a temperate tidal mangrove creek in Botany Bay, New South Wales. *Australian Journal of Freshwater Research*, 35: 33 – 46.
- Bennion, H. and M.A. Smith, 2000. Variability in the water chemistry of shallow ponds in southeast England, with special reference to the seasonality of nutrients and implications for modeling trophic status. *Hydrobiologia*, 436: 145 – 158.
- Berry, P.F., R.P. van der Elst, P. Hanekom, C.S.W. Joubert and M.J. Smale, 1982. Density and biomass of the ichthyofauna of a Natal littoral reef. *Marine Ecology Progress Series*, 10: 49 – 55.
- Blaber, S.J.M. and D.A. Milton, 1990. Species composition, community structure and zoogeography of fishes of mangrove estuaries in the Solomon Islands. *Marine Biology*, 105: 259 – 276.
- Blaber, S.J.M., D.T. Brewer, J.P. Salini, 1989. Species composition and biomasses of fishes in different habitats of a tropical northern Australian estuary: Their occurrence in the adjoining sea and estuarine dependence. *Estuarine, Coastal*

- and Shelf Science, 29: 509 – 531.
- Blaber, S.J.M., D.T. Brewer and A.N. Harris, 1994. Distribution, biomass and community structure of demersal fishes of the Gulf of Carpentaria, Australia. Australian Journal of Marine and Freshwater Research, 45: 375 – 396.
- Blackburn, T.H., 1987. Microbial food webs in sediments. In: Sleigh M.A. (ed.), Microbes in the sea. Ellis Horwood Ltd: New York.
- Blackford, J.C., 1997. An analysis of benthic biological dynamics in a North Sea ecosystem model. Journal of Sea Research, 38(3/4): 213 – 230.
- Blackford, J.C. 1999. The potential role of phytobenthic populations in shallow Northern Adriatic ecosystems. Proceedings of 4th MTP Workshop, Perpignan.
- Blackford, J.C. and P.J. Radford, 1995. A structure and methodology for marine ecosystem modelling. Netherlands Journal of Sea Research, 33(3/4): 247 – 260.
- Briggs, K.T., W.B. Tyler, D.B. Lewis, D.B. and D.R. Carlson, 1987. Bird Communities at Sea off California: 1975-1983. Studies in Avian Biology. 11.
- Bock, M.T., B.S. Miller and A.W. Bowman, 1999. Assessment of eutrophication in the Firth of Clyde: Analysis of coastal water data from 1982 to 1996. Marine Pollution Bulletin, 38: 222 – 231.
- Borkman, D.G. and J.T. Turner, 1993. Plankton studies in Buzzards Bay, Massachusetts, USA II: Nutrients, chlorophyll a and phaeopigments, 1987 to 1990. Marine Ecology Progress Series, 100: 27 – 34.
- Bosserman, R.W., 1980. Complexity measures for assessment of environmental impact in ecosystem networks. In: Proceedings of the Pittsburgh Conference of Modelling and Simulation, Pittsburgh, PA, USA, April 20 – 23, 1980.
- Bosserman, R.W., 1982. Structural comparisons for four lake ecosystem models. In: L. Troncale (ed), Systems Science and Science, Proceedings of the twenty-sixth

- Annual Meeting of the Society for General Systems Research with the American Association for the Advancement of Science, Washington, D.C., January 5 – 9, 1982. Society for General Systems Research: Louisville, Kentucky.
- Botkin, D.B., 1977. Bits, bytes, and IBP. *Bioscience*, 27: 385.
- Boynton, W.R., J.D. Hagy, L. Murray and C. Stokes, 1996. A comparative analysis of eutrophication patterns in a temperate coastal lagoon. *Estuaries*, 19: 408 – 421.
- Breckling, B. and F. Müller, 1994. Current trends in ecological modelling at the 8th ISEM conference on the state-of-the-art. *Ecological Modelling*, 75/76: 667 – 675.
- Brock, I.M., 1977. Trophic relationships in a seagrass community (*Thalassia testudinum*), in Card Sound, Florida: Fish diets in relation to macrobenthic and cryptic faunal abundance. *Transactions of the American Fisheries Society*, 106(3): 219 – 229.
- Broekhuizen, N., 1999. Simulating motile algae using a mixed Eulerian-Lagrangian approach: does motility promote dinoflagellate persistence or co-existence with diatoms? *Journal of Plankton Research*, 21(7): 1191 – 1216.
- Brooks, R.J. and A.M. Tobias, 1996. Choosing the best model: level of detail, complexity, and model performance. *Mathematical Computer Modelling* 24: 1 – 14.
- Broström, G. 1998. A note on the C/N and C/P ratio of the biological production of the Nordic seas. *Tellus*, 50B: 93 – 109.
- Bryant, A.D., M.R. Heath, N. Broekhuizen, J.G. Ollason, W.S.C. Gurney and S.P.R. Greenstreet, 1995. Modelling the predation, growth and population dynamics of fish within a spatially-resolved shelf-sea ecosystem model. *Netherlands Journal*

- of Sea Research, 33: 407 – 421.
- Burford, M.A., P.C. Rothlisberg and Y. Wang, 1995. Spatial and temporal distribution of tropical phytoplankton species and biomass in the Gulf of Carpentaria, Australia. *Marine Ecology Progress Series*, 118: 255 – 266.
- Buskey, E.J., 1994. Annual pattern of micro- and mesozooplankton abundance and biomass in a subtropical estuary. *Journal of Plankton Research*, 15: 907 – 924.
- Büssenschütt, M and C. Pahl-Wostl, 2000. A discrete, allometric approach to the modelling of ecosystem dynamics. *Ecological Modelling*, 126: 33 – 48.
- Butterworth, D.S., 1989. The Benguela Ecology Programme: successful and appropriate? *South African Journal of Marine Science* 85: 633 – 643.
- Butterworth, D.S. and Punt, A.E. 1999. Experiences in the evaluation and implementation of management procedures. *ICES Journal of Marine Science*, 56: 985 – 998.
- Buzzelli, C.P., R.L. Wetzel and M.B. Meyers, 1998. Dynamic simulation of littoral zone habitats in lower Chesapeake bay II: Seagrass habitat primary production and water quality relationships. *Estuaries*, 21: 673 – 689.
- Caddy, J.F., 2000. Marine catchment basin effects versus impacts of fisheries on semi-enclosed seas. *ICES Journal of Marine Science*, 57(3): 628 – 640.
- Cadée, G.C. and J. Hegeman, 1977. Distribution of primary production of the benthic microflora and accumulation of organic matter on a tidal flat area, Balgzand, Dutch Wadden Sea. *Netherlands Journal of Sea Research*, 11: 21 – 41.
- Cale, W.G. and Odell, P.L., 1980. Behaviour of aggregate state variable in ecosystem models. *Mathematical Biosciences*, 49: 121 – 137.
- Cale, W.G. and R.V. O'Neill, 1988. Aggregation and consistency problems in theoretical models of exploitative resource competition. *Ecological Modelling*,

40: 97 – 109.

- Cale, W.G., R.V. O'Neill and R.H. Gardner, 1983. Aggregation error in nonlinear ecological models. *Journal of Theoretical Biology*, 100: 539 – 550.
- Carpenter, S.R. and J.F. Kitchell, 1993. *Aquatic Trophic Cascades*. Cambridge University Press, Cambridge, UK.
- Carrer, S. and S. Opitz, 1999. Trophic network model of a shallow lagoon area in the northern part of the Lagoon of Venice. *Ecological Modelling*, 124: 193 – 219.
- Chavez, E.A., M. Garduno and F. Arreguin-Sanchez, 1993. Trophic dynamic structure of Celestun lagoon, Southern Gulf of Mexico. In V. Christensen and D. Pauly (ed.s) *Trophic Models of aquatic ecosystems*. ICLARM Conference Proceedings, 26: 186 – 192.
- Chidgey, S.S. and M.J. Edmunds, 1997. Standing crop and nutrient content of macrophytes in Port Phillip Bay. Technical Report No. 32. Port Phillip Bay Environmental Study. CSIRO: Canberra, Australia.
- Christensen, V., 1992. Network analysis of trophic interactions in aquatic ecosystems. Royal Danish School of Pharmacy, Copenhagen, Denmark. PhD thesis, pp 55 + 13 Appendices.
- Christensen, V., 1995. Ecosystem maturity – towards quantification. *Ecological Modelling* 77: 3 – 32.
- Christensen, V. and D. Pauly, 1992. ECOPATH II – a software for balancing steady-state ecosystem models and calculating network characteristics. *Ecological Modelling*, 61: 169 – 185.
- Christensen, V., C.J. Walters and D. Pauly 2000. ECOPATH with ECOSIM: A user's guide. October 2000 edition. Fisheries Centre, University of British Columbia, Vancouver, Canada and International Centre for Living Aquatic Resources

Management, Penang, Malaysia.

- Clarke, A. and R.J.G. Leakey, 1996. The seasonal cycle of phytoplankton, macronutrients, and the microbial community in a nearshore Antarctic marine ecosystem. *Limnology and Oceanography*, 41: 1281 – 1294.
- Colijn, F. and V.N. de Jonge, 1984. Primary production of microphytobenthos in the Ems-Dollard estuary. *Marine Ecology Progress Series*, 14: 185 – 196.
- Cohen, J.E. and C.M. Newman, 1988. Dynamic basis of food web organization. *Ecology*, 69: 1655 – 1664.
- Conley, D.J., 1999. Biogeochemical nutrient cycles and nutrient management strategies. *Hydrobiologia*, 410: 87 – 96.
- Connell, J., 1961. The influence of interspecific competition and other factors on the distribution of the barnacle *Chthamalus stellatus*. *Ecology*, 42: 710 – 723.
- Costanza, R and T. Maxwell, 1994. Resolution and predictability: An approach to the scaling problem. *Landscape Ecology*, 9: 47-57.
- Costanza, R. and F.H. Sklar, 1985. Articulation, accuracy and effectiveness of mathematical models: a review of freshwater wetland applications. *Ecological Modelling*, 27: 45 – 68.
- Coutin, P. (ed) 1997. Snapper – 1996. Compiled by the Bay & Inlet Fisheries and Stock Assessment Group. Fisheries Victoria Assessment Report No. 12. Fisheries Victoria: East Melbourne.
- Dauer, D.M. and R.W. Alden III, 1995. Long-term trends in the macrobenthos and water quality of the lower Chesapeake Bay (1985 – 1991). *Marine Pollution Bulletin*, 30: 840 – 850.
- Dean, T.A., L. Haldorson, D.R. Laur, S.C. Jewett, A. Blanchard, 2000. The distribution of nearshore fishes in kelp and eelgrass communities in Prince

- William Sound, Alaska: associations with vegetation and physical habitat characteristics. *Environmental Biology of Fishes*, 57: 271 – 287.
- DeAngelis, D.L., 1975. Stability and connectance in food web models. *Ecology*, 65: 238 – 243.
- DeAngelis, D., L.J. Gross, M.A. Huston, W.F. Wolff, D. M. Fleming, E.J. Comiskey and S.M. Sylvester, 1998. Landscape Modeling for Everglades Ecosystem Restoration. *Ecosystems*, 1: 64 – 75.
- DeCoursey, D.G., 1992. Developing models with more detail: do more algorithms give more truth? *Weed Technology* 6: 709 – 715.
- Del Amo, Y., O. Le Pape, P. Tréguer, B. Quéguiner, A. Ménesguen and A. Aminot, 1997. Impacts of high-nitrate freshwater inputs on macrotidal ecosystems I: Seasonal evolution of nutrient limitation for the diatom-dominated phytoplankton of the Bay of Brest (France). *Marine Ecology Progress Series*, 161: 213 – 224.
- Delgadillo-Hinojosa, F., G. Gaxiola-Castro, J.A. Segovia-Zavala, A. Muñoz-Barbosa and M.V. Orozco-Borbón, 1997. The effect of vertical mixing on primary production in a bay of the Gulf of California. *Estuarine, Coastal and Shelf Science*, 45: 135 – 148.
- Delos Reyes, M. and R. Martens, 1994. Geoecology of Laguna de Bay, Philipines I. Techno-commerical impact on the trophic level structure of the Laguna de Bay aquatic ecosystem 1968 – 1980. *Ecological Modelling*, 75/76: 497 – 509.
- Dickey-Collas, M., R.J. Gowen and C.J. Fox, 1996. Distribution of larval and juvenile fish in the western Irish Sea: relationship to phytoplankton, zooplankton biomass and recurrent physical features. *Marine and Freshwater Research*, 47: 169 – 181.

- Dolphin Research Institute 2000. Dolphin Education Project Question and Answers. Dolphin Research Institute: Frankston, Victoria, Australia. Also available at <http://www.dolphinresearch.org.au/institute.html>.
- Donalson, D.D. and R.M. Nisbet, 1999. Population dynamics and spatial scale: effects of system size on population persistence. *Ecology*, 80: 2492 – 2507.
- Drgas, A., T. Radziejewska and J. Warzocha, 1998. Biomass size spectra of near-shore shallow-water benthic communities in the Gulf of Gdańsk (Southern Baltic Sea). *P.S.Z.N. Marine Ecology*, 19: 209 – 228.
- Droop, M.R., 1973. Some thoughts on nutrient limitation in algae. *Journal of Phycology*, 9: 264 – 272.
- Droop, M.R., 1974. The nutrient status of algal cells in continuous culture. *Journal of the Marine Biology Association of the United Kingdom*, 54: 825 – 855.
- Dunstan, P.K. and C.R. Johnson, submitted. Linking richness, stability and invasion resistance with area in model marine communities.
- Duplisea, D.E., 2000. The ecological hierarchy, model complexity, predictive ability and robust management. In: Report on the Young Scientists Conference on Marine Ecosystem Perspectives, ICES Cooperative Research Report No. 240. International Council for the Exploration of the Sea, Copenhagen, Denmark.
- Duplisea, D.E. and M.V. Bravington, 1999. Harvesting a size-structured ecosystem. ICES 1999/Z:01 Ecosystem Management. International Council for the Exploration of the Sea, Copenhagen, Denmark.
- Durbin, A.G. and E.G. Durbin, 1998. Effects of Menhaden predation on plankton populations in Narragansett Bay, Rhode Island. *Estuaries*, 21: 449 – 465.
- Ebenhöh, W.C., J.G. Baretta-Bekker and J.W. Baretta, 1997. The primary production module in the marine ecosystem model ERSEM II, with emphasis on the light

- forcing. *Journal of Sea Research*, 38: 173 – 194.
- Ebenhöh, W., C. Kohlmeier and P.J. Radford, 1995. The benthic biological submodel in the European Regional Seas Ecosystem Model. *Netherlands Journal of Sea Research*, 33: 423 – 452.
- Edgar, G.J., 1990. The use of size structure of benthic macrofaunal communities to estimate faunal biomass and secondary production. *Journal of Experimental Marine Biology and Ecology*, 137: 195 – 214.
- Edgar, G.J. and C. Shaw, 1995. The production and trophic ecology of shallow-water fish assemblages in southern Australia III: General relationships between sediments, seagrasses, invertebrates and fishes. *Journal of Experimental Marine Biology and Ecology*, 194: 107 – 131.
- Edgar, G.J., C. Shaw, G.F. Watson and L.S. Hammond, 1994. Comparisons of species richness, size-structure and production of benthos in vegetated and unvegetated habitats in Western Port, Victoria. *Journal of Experimental Marine Biology and Ecology*, 176: 201 – 226.
- Edwards, A.M., 2001. Adding detritus to a nutrient-phytoplankton-zooplankton model: a dynamical approach. *Journal of Plankton Research*, 23: 389 – 413.
- Edwards, A.M. and J. Brindley, 1996. Oscillatory behaviour in a three-component plankton population model. *Dynamics and Stability of Systems*, 11: 347 – 370.
- Edwards, A.M. and J. Brindley, 1999. Zooplankton mortality and the dynamical behaviour of plankton population models. *Bulletin of Mathematical Biology*, 61: 303 – 339.
- Edwards, A.M. and A. Yool, 2000. The role of higher predation in plankton population models. *Journal of Plankton Research*, 22: 1085 – 1112.
- Elert, M., A. Butler, J. Chen, C. Dovlete, A. Konoplev, A. Golubenkov, M. Sheppard,

- O. Togawa and T. Zeevaert 1999. Effects of model complexity on uncertainty estimates. *Journal of Environmental Radioactivity*, 42: 255 – 270.
- Elton, C.S., 1958. *Ecology of invasions by animals and plants*. Chapman and Hall: London.
- Fasham, M.J.R., 1993. Modelling the marine biota. In: M. Heimann (ed.), *The Global Carbon Cycle*. Springer-Verlag: Berlin. pp. 457 – 504.
- Finn, J.T., 1976. Measures of ecosystem structure and function derived from analysis of flows. *Journal of Theoretical Biology*, 56: 363 – 380.
- Flores-Verdugo, F., F. González-Farías, F. Amezcua-Linares, A Yáñez-Arancibia, M. Alvarez-Rubio and J.W. Day Jr., 1990. Mangrove ecology, aquatic primary productivity, and fish community dynamics in the Teacapán-Agua Brava lagoon-estuarine system (Mexican Pacific). *Estuaries*, 13: 219 – 230.
- Fogarty, M.J. and S.A. Murawski, 1998. Large-scale disturbance and the structure of marine system-fishery impacts on the Georges Bank. *Ecological Applications*, 8(1 Suppl S): S6 – S22.
- Francis, R.I.C.C., 1992. Recommendations concerning the calculation of maximum constant yield (MCY) and current annual yield (CAY). N.Z. Fisheries Assessment Research Document 92/8.
- Francois, F., J.C. Poggiale, J.P. Durbec and G. Stora, 1997. A new approach for the modelling of the sediment reworking induced by a macrobenthic community. *Acta Biotheretica*, 45: 295 – 319.
- Francour, P., 1997. Fish assemblages of *Posidonia oceanica* beds at Port-Cros (France, NW Mediterranean): Assessment of composition and long-term fluctuations by visual census. *P.S.Z.N. Marine Ecology*, 18(2): 157 – 173.

- Fransz, H.G., Mommaerts, J.P. and G. Radach, 1991. Ecological Modelling of the North Sea. Netherlands Journal of Sea Research, 28(1/2): 67-140.
- Fredette, T.J., R.J. Diaz, J. van Montfrans and R.J. Orth, 1990. Secondary production within a seagrass bed (*Zostera marina* and *Ruppia maritima*) in Lower Chesapeake Bay. Estuaries, 13: 431 – 440.
- Freidlander, A.M. and J.D. Parrish, 1997. Fisheries harvest and standing stock in a Hawaiian bay. Fisheries Research, 32: 33 – 50.
- Frost, B.W., 1975. A threshold feeding behaviour in *Calanus pacificus*. Limnology and Oceanography, 20: 263 – 266.
- Gallegos, C.L., T.E. Jordan and D.L. Correll, 1997. Interannual variability in spring bloom timing and magnitude in the Rhode River, Maryland, USA: observations and modelling. Marine Ecology Progress Series, 154: 27 – 40.
- Gao, H, H. Wei, W. Sun and X. Zhai, 2000. Functions used in biological models and their influences on simulations. Indian Journal of Marine Science, 29: 230 – 237.
- Gardner, M.R. and W.R. Ashby, 1970. Connectance of large dynamic (cybernetic) systems: critical values for stability. Nature, 228: 784.
- Gardner, R.H., W.G. Cale and R.V. O'Neill, 1982. Robust analysis of aggregation error. Ecology, 63: 1771 – 1779.
- Gasca, R. and I. Castellanos, 1993. Zooplankton of Chetumal Bay, Caribbean of Mexico. Revista de Biología Tropical, 41: 619 – 625.
- Gislason, H., M. Sinclair, K. Sainsbury and R. O'Boyle, 2000. Symposium overview: incorporating ecosystem objectives within fisheries management. ICES Journal of Marine Science, 57: 468 – 475.
- Gerral, M and M. Verlaque, 1995. Soft-bottom macrophytobenthos and associated

- environmental factors in Thau lagoon (France, Mediterranean). *Oceanologica Acta*, 18(5): 557 – 571.
- Gibbs, M.M. and W.N. Vant, 1997. Seasonal changes in factors controlling phytoplankton growth in Beatrix Bay, New Zealand. *N.Z. Journal of Marine and Freshwater Research*, 31: 237 – 248.
- Gibson, R.N. and A.D. Ansell and L. Robb, 1993. Seasonal and annual variations in abundance and species composition of fish and macrocrustacean communities on a Scottish sandy beach. *Marine Ecology Progress Series*, 98: 89 – 105.
- Ginzburg, L.R. and H.R. Akçakaya, 1992. Consequences of ratio-dependent predation for steady-state properties of ecosystems. *Ecology*, 73: 1536 – 1543.
- Gleeson, S.K., 1994. Density dependence is better than ratio dependence. *Ecology*, 75: 1834 – 1835.
- Goswami, S.C. and G. Padmavati, 1996. Zooplankton production, composition and diversity in the coastal waters of Goa. *Canadian Journal of Fisheries and Aquatic Sciences*, 25: 91 – 97.
- Gray, J.S., 1992. Eutrophication in the sea. In: G. Colombo, I. Ferrari, V. Ceccherelli and R. Rossi (ed.s), *Marine Eutrophication and Population Dynamics: 25th European Marine Biology Symposium*, University of Ferrara, pp: 3 – 15.
- Grémare, A., J.M. Amouroux and G. Vétion, 1998. Long-term comparison of macrobenthos within the soft bottoms of the Bay of Banyuls-sur-mer (northwestern Mediterranean Sea). *Journal of Sea Research*, 40: 281 – 302.
- Grøntved, J., 1960. On the productivity of microbenthos and phytoplankton in some Danish fjords. *Meddelelser Fra Danmarks Fiskeri – Og Havundersøgelser. N.S.*, 3: 55 – 92.
- Grove, S.L. and P.K. Probert, 1999. Sediment macrobenthos of upper Otago Harbour,

- New Zealand. N.Z. Journal of Marine and Freshwater Research, 33: 469 – 480.
- Guillaud, J.F., F. Andrieux and A. Menesguen, 2000. Biogeochemical modelling in the Bay of Seine (France): an improvement by introducing phosphorus in nutrient cycles. Journal of Marine Systems, 25: 369 – 386.
- Gunthorpe, L., P. Hamer and S. Walker, 1997. Bays and Inlets Scalefish Fisheries Review: Volume 1: Life cycles and habitat requirements of selected Victorian fish species, including an assessment of the main habitat threatening processes and recommendations for habitat maintenance, rehabilitation and enhancement. Marine and Freshwater Resources Institute: Queenscliff, Victoria, Australia.
- Gurney, W.S.C and R.M. Nisbet, 1978. Predator-prey fluctuations in patchy environments. Journal of Animal Ecology, 47: 85 – 102.
- Hagen, N.T., 1995. Recurrent destructive grazing of successional immature kelp forests by Green Sea Urchins in Vestfjorden, Northern Norway. Marine Ecology Progress Series, 123: 95 – 106.
- Hairstone, N.G., F.E. Smith and L.B. Slobodkin, 1960. Community structure, population control and competition. American Naturalist, 94: 421 – 425.
- Halfon, E., 1979. Computer-based development of large-scale ecological models: problems and prospects. In: G.S. Innis and R.V. O'Neill (eds), Systems analysis of ecosystems. International Co-operative Publishing House: Fairland, Maryland.
- Halfon, E., 1983a. Is there a best model structure? I. Modeling the fate of a toxic substance in a lake. Ecological Modelling, 20: 135-152.
- Halfon, E., 1983b. Is there a best model structure? II. Comparing the Model structures of different fate models. Ecological Modelling, 20: 153-163.
- Hall, D.N. (ed), 1992. Port Phillip Bay Environmental Study: Status Review.

- Technical Report No. 9. Port Phillip Bay Environmental Study. CSIRO:
Canberra, Australia.
- Hall, J.A., D.P. Barrett and M.R. James, 1993. The importance of phytoflagellate, heterotrophic flagellate and ciliate grazing on bacteria and picophytoplankton sized prey in a coastal marine environment. *Journal of Plankton Research*, 15: 1075 – 1086.
- Håkanson, L., 1995. Optimal size of predictive models. *Ecological Modelling*, 78:195-204
- Håkanson, L., 1997. Modelling of Radiocesium in Lakes – on Predictive Power and Lessons for the Future. In: G. Desmet, R.J.Blust, R.N.J.Comans, J.A.Fernandez, J.Hilton and A.Bettencourt (eds), *Freshwater and Estuarine Radioecology*, Studies in Environmental Science 68.
- Harding, S.P., 1999. Food web complexity enhances community stability and climate regulation in a geophysical model. *Tellus*, 51B: 815 – 829.
- Hargrave, B.T., N.J. Prouse, G.A. Phillips and P.A. Neame, 1983. Primary production and respiration in pelagic and benthic communities at two intertidal sites in the upper bay of Fundy. *Canadian Journal of Fisheries and Aquatic Sciences*, 40(Suppl. 1): 229 – 243.
- Harris, G., G. Batley, D. Fox, D. Hall, P. Jernakoff, R. Molloy, A. Murray, B. Newell, J. Parslow, G. Skyring and S. Walker, 1996. Port Phillip Bay Environmental Study Final Report. CSIRO, Canberra, Australia.
- Harvey, W.H., 1950. On the production of living matter in the sea off Plymouth. *J. Mar. Biol. Assoc.* 24: 97 – 137.
- Hassell, M.P., 1976. The dynamics of competition and predation. Edward Arnold: London.

- Hassell, M.P. and H.N. Commins, 1978. Sigmoid functional responses and population stability. *Theoretical Population Biology*, 14: 62 – 67.
- Hassell, M.P. and R.M. May, 1973. Stability in insect host-parasite models. *Journal of Animal Ecology*, 42: 693 – 736.
- Hassell, M.P., H.N. Comins and R. May, 1994. Species coexistence and self-organizing spatial dynamics. *Nature*, 370: 290 – 292.
- Heck, K.L. Jr., K.W. Able, C.T. Roman and M.P. Fahay, 1995. Composition, abundance, biomass, and production of macrofauna in a New England estuary: Comparisons among eelgrass meadows and other nursery habitats. *Estuaries*, 18: 379 – 389.
- Hedgpeth, J.W., 1977. Models and muddles: some philosophical observations. *Helgoländer wiss. Meeresunters*, 30: 92 – 104.
- Herbert, R.A., 1999. Nitrogen cycling in coastal marine ecosystems. *FEMS Microbiology Reviews*, 23(5): 563 – 590.
- Heymans, J.J. and D. Baird, 1995. Energy flow in the Kromme estuarine ecosystem, St Francis Bay, South Africa. *Estuarine, Coastal and Shelf Science*, 41: 39 – 59.
- Hibbert, C.J., 1976. Biomass and production of a bivalve community on an intertidal mud-flat. *Journal of Experimental Marine Biology and Ecology*, 25: 249 – 261.
- Hoch, T. and P. Garreau, 1998. Phytoplankton dynamics in the English Channel: a simplified three-dimensional approach. *Journal of Marine Systems*, 16: 133 – 150.
- Holling, C.S., 1959. Some characteristics of simple types of predation and parasitism. *Canadian Entomologist* 91: 385 – 398.

- Holloway, M., and G. Jenkins, 1993. The Role of Zooplankton in Nitrogen and Carbon Cycling in Port Phillip Bay. Technical Report No. 11. Port Phillip Bay Environmental Study. CSIRO: Canberra, Australia.
- Hollowed, A.B., N. Bax, R. Beamish, J. Collie, M. Fogarty, P., Livingston, J. Pope and J.C. Rice, 2000. Are multispecies models an improvement on single-species models for measuring fishing impacts on marine ecosystems? ICES Journal of Marine Science, 57: 707 – 719.
- Holmes, EE., M.A. Lewis, J.E. Banks and R.R. Veit, 1994. Partial differential equations in ecology – spatial interactions and population dynamics. Ecology, 75(1): 17 – 29.
- Hopcroft, R.R., J.C. Roff and D. Lombard, 1998. Production of tropical copepods in Kingston Harbour, Jamaica: the importance of small species. Marine Biology, 130: 593 – 601.
- Houde, E.D. and E.S. Rutherford, 1993. Recent trends in estuarine fisheries – predictions of fish production and yield. Estuaries, 16(2): 161 – 176.
- Huffaker, C.B., 1958. Experimental studies on predation: dispersion factors and predator-prey oscillations. Hilgardia, 27: 343 – 383.
- Hurt G.C and R.A. Armstrong, 1996. A pelagic ecosystem model calibrated with BATS data. Deep-Sea Research II, 43: 653-683.
- Innis, G.S. and E. Rexstad, 1983. Simulation and model simplification techniques. Simulation, 41: 7 – 15.
- Iriarte, A., I. de Madariaga, F. Deiz-Garagarza, M. Revilla and E. Orive, 1996. Primary plankton production, respiration and nitrification in a shallow temperate estuary during summer. Journal of Experimental Marine Biology and Ecology, 208: 127 – 151.

- Iwasa, Y., V. Andreassen and S. Levin, 1987. Aggregation in model ecosystems I: Perfect aggregation. *Ecological Modelling*, 37: 287 – 302.
- Jacobson, D.M., 1999. A brief history of dinoflagellate feeding research. *Journal of Eukaryotic Microbiology*, 46: 376 – 381.
- Jean, F. and G. Thouzeau, 1995. Estimate of state variables of a trophic web model in the bay of Brest. *Comptes Rendus de l'Academie des Sciences Serie III – Sciences de la Vie-Life Sciences*. 318: 145 – 154.
- Jeong, H.J., J.H. Shim, J.S. Kim, J.Y. Park, C.W. Lee and Y. Lee, 1999. Feeding by the mixotrophic thecate dinoflagellate *Fragilidium cf. Mexicanum* on red-tide and toxic dinoflagellates. *Marine Ecology Progress Series*, 176: 263 – 277.
- Jester, D.B. Jr, D.L. Jr Garling, A.R. Tipton and R.T. Lackey, 1977. A general population dynamics theory for Largemouth Bass. FWS 1 – 77, Virginia Polytechnic Inst. and State Univ.
- Johnson, C.R., 1997. Self-organizing in spatial competition systems. In: N.I. Klomp and I.D. Lunt (eds), *Frontiers in ecology – building the links*. Elsevier: Amsterdam.
- Jørgensen, S.E., 1994. *Fundamentals of ecological modelling*, 2nd ed. *Developments in Environmental Modelling* 19. Elsevier: Amsterdam, London, New York.
- Jørgensen, S.E., B.C. Patten and M. Straškraka, 1992. Ecosystems emerging: toward an ecology of complex systems in a complex future. *Ecological Modelling*, 62: 1 – 27.
- Kailola, P.J., M.J. Williams, P.C. Stewart, R.E. Reichelt, A. McNee and C. Grieve, 1993. *Australian Fisheries Resources*. Bureau of Resource Sciences, Department of primary Industries and Energy and the Fisheries Research and Development Corporation: Canberra, Australia.

- Kallianiotis, A., K. Sophronidis, P. Vidoris and A. Tselepides, 2000. Demersal fish and megafaunal assemblages on the Cretan continental shelf and slope (NE Mediterranean): seasonal variation in species density, biomass and diversity. *Progress in Oceanography*, 46: 429 – 455.
- Keitt, T.H. 1997. Stability and complexity on a lattice: coexistence of species in an individual-based food web model. *Ecological Modelling*, 102: 243 – 258.
- Kinney, E.H and C.T. Roman, 1998. Response of primary producers to nutrient enrichment in a shallow estuary. *Marine Ecology Progress Series*, 163: 89 – 98.
- Kjørboe, T. and T.G. Nielsen, 1994a. Regulation of zooplankton biomass and production in a temperate, coastal ecosystem 1: Copepods. *Limnology and Oceanography*, 39: 493 – 507.
- Kjørboe, T. and T.G. Nielsen, 1994b. Regulation of zooplankton biomass and production in a temperate, coastal ecosystem 2: Ciliates. *Limnology and Oceanography*, 39: 493 – 507.
- Kjerfve, B., C.H.A. Rjbeiro, G.T.M. Dias, A.M. Filippo and V. Da Silva Quaresma, 1997. Oceanographic characteristics of an impacted coastal bay: Baía de Guanabara, Rio de Janeiro, Brazil. *Continental Shelf Research*, 17: 1609 – 1643.
- Koopman, M.T., A.E. Punt and D.C. Smith, 2000. Production parameters from the fisheries literature for SEF-like species. Final Report to Australian Fisheries Management, ARF Project R99/0308.
- Koski, M., M. Viitasalo and H. Kuosa, 1999. Seasonal development of meso-zooplankton biomass and production on the SW coast of Finland. *Ophelia*, 50: 69 – 91.
- Kremer J.N. and P. Kremer, 1982. A three-trophic level estuarine model: Synergism

- of two mechanistic simulations. *Ecological Modelling*, 415: 145-157.
- Kuiter, R.H., 1996. Guide to sea fishes of Australia. New Holland Publishers Ltd: London, Sydney.
- Kulbicki, M., P. Labrosse and Y. Letourneur, 2000. Fish stock assessment of the northern New Caledonian lagoons: 2 – Stocks of lagoon bottom and reef-associated fishes. *Aquatic Living Resources*, 13(2): 77 – 90.
- Kumari, L.K. and S.C. Goswami, 1993. Biomass and biochemical composition of zooplankton from northwest Bay of Bengal during January 1990. *Indian Journal of Marine Science*, 22: 143 – 145.
- Leach, J.H., 1970. Epibenthic algal production in an intertidal mudflat. *Limnology and Oceanography*, 15: 514 – 521.
- Lee, P.G., 1994. Nutrition of Cephalopods: Fueling the System. *Marine Behaviour and Physiology*, 25 (1-3): 35-51.
- Lee, K. and P.A. Fishwick, 1998. Generation and selection of the optimal model for real-time simulation. SPIE Aerosense Conference, April, 1998, Orlando, Florida. pp. 164 – 175.
- Legrand, C., E. Graneli and P. Carlsson, 1998. Induced phagotrophy in the photosynthetic dinoflagellate *Heterocapsa triquetra*. *Aquatic Microbial Ecology*, 15: 65 – 75.
- Leibold, M.A. 1996. A graphical model of keystone predators in food webs: trophic regulation of abundance, incidence, and diversity patterns in communities. *American Naturalist*, 147: 784 – 812.
- Leinaas, H.P. and H. Christie, 1996. Effects of removing sea urchins (*Strongylocentrotus droebachiensis*) – stability of the barrens state and succession of kelp forest recovery in the east Atlantic. *Oecologia*, 105: 524 –

- Letourneur, Y., 1998. Composition, structures et réseaux trophiques des peuplements de poissons de la côte au vent de l'île de La Réunion. *Cybiurn*, 22: 267 – 283.
- Letourneur, Y., M. Kulbicki and P. Labrosse, 2000. Fish stock assessment of the northern New Caledonian lagoons: 1 – Structure and stocks of coral reef fish communities. *Aquatic Living Resources*, 13(2): 65 – 76.
- Levin, S.A., 1992. The problem of pattern and scale in ecology. *Ecology*, 73: 1943 – 1967.
- Levins, R., 1970. Extinction. In: M. Gerstenhaber (ed), *Some Mathematical Problems in Biology. Lectures on Mathematics in Life Sciences Vol II*. The American Mathematical Society: Providence. Pp 77 – 107.
- Levinton, J.S., 1995. *Marine Biology: Function, biodiversity, ecology*. Oxford University Press: New York, Oxford.
- Li, A.S., D.K. Stoecker and J.E. Adolf, 1999. Feeding, pigmentation, photosynthesis and growth of the mixotrophic dinoflagellate *Gyrodinium galatheanum*. *Aquatic Microbial Ecology*, 19: 163 – 176.
- Li, A.S., D.K. Stoecker and D.W. Coats, 2000. Mixotrophy in *Gyrodinium galatheanum* (Dinophyceae): Grazing responses to light intensity and inorganic nutrients. *Journal of Phycology*, 36: 33 – 45.
- Lima, I.D. and J.P. Castello, 1995. Distribution and abundance of South-west Atlantic anchovy spawners (*Engraulis anchoita*) in relation to oceanographic processes in the Southern Brazilian Shelf. *Fisheries Oceanography*, 4(1): 1 – 16.
- Lin, H.J., K.T. Shao, S.R. Kuo, H.L. Hsieh, S.L. Wong, L.M. Chen, W.T. Lo and J.J. Hung, 1999. A trophic model of a sandy barrier lagoon at Chiku in southwestern Taiwan. *Estuarine, Coastal and Shelf Science*, 48: 575 – 588.

- Loneragan, N.R., I.C. Potter, R.J.C. Lenanton and N. Caputi, 1986. Spatial and seasonal differences in the fish fauna in the shallows of a large Australian estuary. *Marine Biology*, 92: 575 – 586.
- Lotze, H.K., W. Schramm, D. Schories and B. Worm, 1999. Control of macroalgal blooms at early developmental stages: *Pilayella littoralis* versus *Enteromorpha* spp. *Oecologia*, 119(1): 46 – 54.
- Love M.S., J.E. Caselle and L. Snook, 2000. Fish assemblages around seven oil platforms in the Santa Barbara Channel area. *Fisheries Bulletin*, 98: 96 – 117.
- Lubbers, L., W.R. Boynton and W.M. Kemp, 1990. Variations in structure of estuarine fish communities in relation to abundance of submersed vascular plants. *Marine Ecology Progress Series*, 65: 1 – 14.
- Ludwig, D. and C.J. Walters, 1981. Measurement errors and uncertainty in parameter estimates for stock recruitment. *Canadian Journal of Fisheries and Aquatic Sciences*, 38: 711-720.
- Ludwig, D. and C.J. Walters, 1985. Are age-structured models appropriate for catch-effort data? *Canadian Journal of Fisheries and Aquatic Sciences*, 42: 1066 – 1072.
- MacArthur, R.H., 1955. Fluctuations of animal populations and a measure of community stability. *Ecology*, 36: 533 – 536.
- MacArthur, R.H. and E.O. Wilson, 1967. *The Theory of Island Biogeography*. Princeton University Press: Princeton.
- MacIntyre, H.L., R.J. Geider and D.C. Miller, 1996. Microphytobenthos: The ecological role of the “secret garden” of unvegetated, shallow-water marine habitats I: Distribution, abundance and primary production. *Estuaries*, 19: 186 – 201.

- Mace, P.M., 2001. A new role for MSY in single-species and ecosystem approaches to fisheries stock assessment and management. *Fish and Fisheries*, 2: 2 – 32.
- Madden, C.J. and W.M. Kemp, 1996. Ecosystem model of an estuarine submersed plant community: calibration and simulation of eutrophication responses. *Estuaries*, 19: 457 – 474.
- Manickchand-Heileman, S., F. Arreguín-Sánchez, A. Lara-Domínguez and L.A. Soto, 1998. Energy flow and network analysis of Terminos Lagoon SW Gulf of Mexico. *Journal of Fish Biology*, 53(Supl. A): 179 – 197.
- Matheke, G.E.M and R. Horner, 1974. Primary productivity of the benthic microalgae in the Chukchi Sea near Barrow, Alaska. *Journal of the Fisheries Research Board of Canada*, 31: 1779 – 1786.
- May, R.M., 1973. *Stability and complexity in model ecosystems*. Princeton University Press: Princeton, New Jersey.
- May, R.M., 1974. *Stability and Complexity in Model Ecosystems*, 2nd edn. Monographs in Population Biology No. 6. Princeton University Press: Princeton.
- May, R.M., 1976. *Theoretical ecology*. Saunders: Philadelphia.
- Maynard Smith, J., 1974. *Models in Ecology*. Cambridge University Press: Cambridge.
- Matson, P. and A.A. Berryman, 1992. Special feature: ratio-dependent predator-prey theory. *Ecology*, 73: 1529 – 1566.
- McCann, K.S., 2000. The diversity-stability debate. *Nature*, 405: 228 – 233.
- McCann, K., A. Hastings and G.R. Huxel, 1998. Weak trophic interactions and the balance of nature. *Nature*, 395: 794 – 798.
- McClanahan, T.R. and E. Sala, 1997. A Mediterranean rocky-bottom ecosystem

- fisheries model. *Ecological Modelling*, 104: 145 – 164.
- McCreary, J.P. Jr., Kohler, K.E., Hood, R.R. and D.B. Olson, 1996. A Four-Component Ecosystem Model of Biological Activity in the Arabian Sea. *Progress in Oceanography*, 37: 193 – 240.
- McGillicuddy, D.J., J.J. McCarthy and A.R. Robinson, 1995. Coupled physical and biological modelling of the spring bloom in the north Atlantic (I): Model formulation and one dimensional bloom processes. *Deep-Sea Research*, 42: 1313 – 1357.
- Menasveta, P., T. Wongratana, N. Chaitanawisuti and S. Rungsupa, 1986. Species composition and standing crop of coral reef fishes in Sichang Islands, Gulf of Thailand. *Galaxea*, 5: 115 – 121.
- Mills, E.L., 1975. Benthic organisms and the structure of marine ecosystems. *Journal of the Fisheries Research Board of Canada*, 32: 1657 – 1663.
- Möller, P. and R. Rosenberg, 1982. Production and abundance of the amphipod *Corophium volutator* on the west coast of Sweden. *Netherlands Journal of Sea Research*, 16: 127 – 140.
- Möller, P., L. Pihl and R. Rosenberg, 1985. Benthic faunal energy flow and biological interaction in some shallow marine soft bottom habitats. *Marine Ecology Progress Series*, 27: 109 – 121.
- Monbet, Y., 1992. Control of phytoplankton biomass in estuaries: A comparative analysis of microtidal and macrotidal estuaries. *Estuaries*, 15: 563 – 571.
- Morris, M.D., 1991. Factorial sampling plans for preliminary computational experiments. *Technometrics*, 33(2): 161 – 174.
- Mortazavi, B., R.L. Iverson, W.M. Landing, F.G. Lewis and W. Huang, 2000. Control of phytoplankton production and biomass in a river-dominated estuary:

- Apalachicola Bay, Florida, USA. Marine Ecology Progress Series, 198: 19 – 31.
- Morton, R.M., 1990. Community structure, density and standing crop of fishes in a subtropical Australian mangrove area. Marine Biology, 105: 385 – 394.
- Murray, A.G., 2001. The use of simple models in the design and calibration of a dynamic 2D model of a semi-enclosed Australian bay. Ecological Modelling, 136: 15 – 30.
- Murray, A., and J. Parslow, 1997. Port Phillip Bay Integrated Model: Final Report. Technical Report No. 44. Port Phillip Bay Environmental Study. CSIRO: Canberra, Australia.
- Murray, A.G. and J.S. Parslow, 1999a. Modelling of nutrient impacts in Port Phillip Bay – a semi-enclosed marine Australian ecosystem. Marine and Freshwater Research, 50(6): 597 – 611.
- Murray, A.G. and J.S. Parslow, 1999b. The analysis of alternative formulations in a simple model of a coastal ecosystem. Ecological Modelling, 119: 149 – 166.
- Murray L, and R.L. Wetzel, 1987. Oxygen production and consumption associated with the major autotrophic components in two temperate seagrass communities. Marine Ecology Progress Series, 38: 231 – 239.
- Nicholson, G.J., A.R. Longmore and R.A. Cowdell, 1996. Nutrient Status of the Sediments of Port Phillip Bay. Technical Report No. 26. Port Phillip Bay Environmental Study. CSIRO: Canberra, Australia.
- Nielsen, E. and K. Richardson, 1996. Can changes in the fisheries yield in the Kattegat (1950 – 1992) be linked to changes in primary production. ICES Journal of Marine Science, 53(6): 988 – 994.
- Nienhuis, P.H., 1983. Temporal and spatial patterns of eelgrass (*Zostera marina* L.) in a former estuary of the Netherlands, dominated by human activities. Marine

- Technology Society Journal, 17: 69 – 77.
- Nihoul, J.C.J., 1998a. Optimum complexity in ecohydrodynamic modelling: An ecosystem dynamics standpoint. *Journal of Marine Systems*, 16: 3 – 5.
- Nihoul, J.C.J., 1998b. Modelling marine ecosystems as a discipline in Earth Science. *Earth-Science Review*, 44: 1 – 13.
- Nihoul, J.C.J and E.J.M. Delhez, 1998. Preface, 28th International Liège Colloquium on Ocean Hydrodynamics: Liège, May 6 – 10, 1996: Modelling hydrodynamically dominated marine ecosystems. *Journal of Marine Systems*, 16: 1 – 2.
- Nihoul, J.C.J. and S. Djenidi, 1998. Chapter 18: Coupled physical, chemical and biological models. In: K.H. Brink and A.R. Robinson (eds), *The Sea*. John Wiley & Sons Inc: New York, London.
- Nisbet, R.M. and W.S.C Gurney, 1976. A simple mechanism for population cycles. *Nature*, 334: 147 – 149.
- Odum, E.P., 1953. *Fundamentals of ecology*. Saunders: Philadelphia
- Odum, H.T. and E.P. Odum, 1955. Trophic structure and productivity of a windward coral reef community on Eniwetok Atoll. *Ecological Monographs*, 25: 291 – 320.
- Officer, R.A., and G.D. Parry, 1996. *Food Webs of Demersal Fish in Port Phillip Bay*. Technical Report No. 36. Port Phillip Bay Environmental Study. CSIRO: Canberra, Australia.
- Olivieri, R.A. and F.P. Chavez, 2000. A model of plankton dynamics for the coastal upwelling system of Monterey Bay, California. *Deep-Sea Research II*, 47: 1077 – 1106.

- Olson, D.B. and R.R. Hood, 1994. Modelling Pelagic Biogeography. *Progress in Oceanography*, 34: 161-205.
- O'Neill, R.V., 1975. Modeling in the eastern deciduous forest biome. In: B.C. Patten (ed), *System analysis and simulation in ecology*, III. Academic Press: New York, London. pp: 49 – 72.
- O'Neill, R.V., 1979. Transmutations across hierarchical levels. In: G.S. Innis and R.V. O'Neill (eds.), *Systems analysis of ecosystems*. International Co-operative: Fairland, Maryland USA. pp: 59 – 78.
- O'Neill, R.V. and B. Rust, 1979. Aggregation error in ecological models. *Ecological Modelling*, 7: 91 – 105.
- Optiz, S., 1996. Trophic interactions in Caribbean coral reefs. ICLARM Technical Report 43.
- Osterroht, C. and H. Thomas, 2000. New production enhanced by nutrient supply from non-Redfield remineralisation of freshly produced organic material. *Journal of Marine Systems*, 25: 33 – 46.
- Pahl-Wostl, C., 1997. Dynamic structure of a food web model: comparison with a food chain model. *Ecological Modelling*, 100: 103 – 123.
- Paine, R.T., 1966. Food web complexity and species diversity. *American Naturalist*, 100: 65 – 75.
- Palomares, M.L.D., P. Reyes-Marchant, N. Lair, M. Zainure, G. Barnabé and G. Lasserre, 1993. A trophic model of a Mediterranean Lagoon, Etang de Thau, France. In V. Christensen and D. Pauly (ed.s) *Trophic Models of aquatic ecosystems*. ICLARM Conference Proceedings, 26: 20 – 28.
- Park, G.S. and H.G. Marshal, 2000. Estuarine relationships between zooplankton community structure and trophic gradients. *Journal of Plankton Research*, 22:

121 – 135.

- Parry, G.D., D.K. Hobday, D.R. Currie, R.A. Officer and A.S. Gason, 1995. The Distribution, Abundance and Diets of Demersal Fish in Port Phillip Bay. Technical Report No. 21. Port Phillip Bay Environmental Study. CSIRO: Canberra, Australia.
- Parsons, T.R. and Y.L.L. Chen, 1994. Estimates of trophic efficiency, based on the size distribution of phytoplankton and fish in different environments. *Zoological Studies*, 33(4): 296 – 301.
- Pascual, M. and S. A. Levin, 1999. From individuals to population densities: searching for the intermediate scale of nontrivial determinism. *Ecology*, 80: 2225 – 2236.
- Patten, B.C, D.A. Egloff and T.H. Richardson, 1975. Total ecosystem model for a cove in Lake Texoma. In: B.C. Patten (ed), *Systems analysis and simulation in ecology*, vol III. Academic Press: New York.
- Pennington, J.T. and F.P. Chavez, 2000. Seasonal fluctuations of temperature, salinity, nitrate, chlorophyll and primary production at station H3/M1 over 1989 – 1996 in Monterey Bay, California. *Deep-Sea Research II*, 47: 947 – 973.
- PICES Working Group 11, 1998. Consumption of Marine Resources by Marine Birds and Mammals in the PICES Region. 1998 Report of Working Group 11. Sidney, Canada. Also available at <http://pices.ios.bc.ca/wg/wgf.htm>.
- Pihl, L., H. Wennhage and S. Nilsson, 1994. Fish assemblage structure in relation to macrophytes and filamentous epiphytes in shallow non-tidal rocky- and soft-bottom habitats. *Environmental Biology of Fishes*, 39: 271 – 288.
- Pihl-Baden, S. and L. Pihl, 1984. Abundance, biomass and production of motile epibenthic fauna in *Zostera marina* (L.) meadows, western Sweden. *Ophelia* 23:

65 – 90.

Pimm, S.L., 1982. Food Webs. Chapman and Hall: London.

Pimm, S.L. and J.H. Lawton, 1978. On feeding on more than one trophic level.

Nature, 275: 542 – 544.

Platt, T.K., H. Mann and R.E. Ulanowicz (eds.), 1981. Mathematical Models in

Biological Oceanography. The UNESCO Press:Paris. 156p.

Plounevez, S. and G. Champalbert, 1999. Feeding behaviour and trophic environment

of *Engraulis encrasicolus* (L.) in the Bay of Biscay. Estuarine, Coastal and

Shelf Science, 49: 177 – 191.

Polis, G.A. and D.R. Strong, 1996. Food web complexity and community dynamics.

American Naturalist, 147: 813 – 846.

Poore, G.C.B, 1992. Soft-bottom macrobenthos of Port Phillip Bay: a literature

review. Technical Report No. 2. Port Phillip Bay Environmental Study. CSIRO:

Canberra, Australia.

Poulet, S.A., M. Laabir and Y. Chaudron, 1996. Characteristic features of

zooplankton in the Bay of Biscay. Scientia Marina, 60(Supl. 2): 79 – 95.

Power, D. and D.B. Atkinson, 1998. The status of Redfish in Unit 2 (Laurentian

Channel Management Unit). DFO Canadian Stock Assessment Secretariat

Research Document 98/021. Canadian Stock Assessment Secretariat, Fisheries

and Oceans (Station 1256), Ottawa, Canada.

Punt, A.E., A.D.M. Smith and G. Cui. (In press) Evaluation of management tools for

Australia's South East Fishery. 2. How well do commonly-used stock

assessments methods perform?. Marine and Freshwater Research.

Putman, R.J., 1994. Community ecology. Chapman and Hall: New York, London. pg

44.

- Quast, J.C., 1968. Estimates of the populations and the standing crop of fishes. *Fish Bulletin*, 139: 57 – 79.
- Radach, J. and A. Moll, 1993. Estimation of the variability of production by simulating annual cycles of phytoplankton in the central North Sea. *Progress in Oceanography*, 31: 339 – 419.
- Radford 1996. The Modules of the European Regional Seas Ecosystem Model II, 1993 – 1996. Plymouth Marine Laboratory, Plymouth, UK.
- Rantatajärvi, E., R. Olsonen, S. Hällfors, J.-M. Leppänen and M. Raateoja, 1998. Effect of sampling frequency on detection of natural variability in phytoplankton: unattended high-frequency measurements on board ferries in the Baltic Sea. *ICES Journal of Marine Science*, 55: 697 – 704.
- Rastetter, E.B., A.W. King, B.J. Cosby, G.H. Hornberger, R.V. O'Neill and J.H. Hobbie, 1992. Aggregating fine-scale ecological knowledge to model coarser-scale attributes of ecosystems. *Ecological Applications*, 2: 55 – 70.
- Redfield, A.C., B.H. Ketchum and F.A. Richards 1963. The influence of organisms on the composition of seawater. In: M.N. Hill (ed), *The sea volume 2*. Wiley-Interscience: New York.
- Reichert, P. and M. Omlin, 1997. On the usefulness of overparameterized ecological models. *Ecological Modelling*, 95: 289 – 299.
- Reigman, R. and B.R. Kuipers, 1993. Resource Competition and Selective Grazing of Plankton in a Multispecies Pelagic Food Web Model. *Marine Ecology*, 15(2): 153-164.
- Rhodes, K.L., 1998. Seasonal trends in epibenthic fish assemblages in the near-shore waters of the western Yellow Sea, Qingdao, People's Republic of China. *Estuarine, Coastal and Shelf Science*, 46: 629 – 643.

- Rhodes, O.E. and R. K. Chesser (eds), 1996. Population Dynamics in Ecological Space and Time. University of Chicago Press: London.
- Robertson, A.I., 1980. The structure and organization of an eelgrass fish fauna. *Oecologia*, 47: 76 – 82.
- Robertson, A.I., 1984. Trophic interactions between the fish fauna and macrobenthos of an eelgrass community in Western Port, Victoria. *Aquatic Botany*, 18: 135 – 153.
- Robertson, A.I. and N.C. Duke, 1990. Mangrove fish-communities in tropical Queensland, Australia: spatial and temporal patterns in densities, biomass and community structure. *Marine Biology*, 104: 369 – 379.
- Roden, E.E. and J.H. Tuttle, 1996. Carbon cycling in mesohaline Chesapeake Bay sediment: Kinetics of particulate and dissolved organic carbon turnover. *Journal of Marine Research*, 54: 343 – 383.
- Rosado-Solorzano, R. and S.A. Guzmán del Prío, 1998. Preliminary trophic structure model of Tampamachoo lagoon, Veracruz, Mexico. *Ecological Modelling*, 109: 141 – 154.
- Ross, S.T., R.H. McMichael Jr, and D.L. Rupel, 1987. Seasonal and diel variation in the standing crop of fishes and macroinvertebrates from a Gulf of Mexico surf zone. *Estuarine, Coastal and Shelf Science*, 25: 391 – 412.
- Roughgarden, J., S. Gaines and H. Possingham, 1988. Recruitment dynamics in complex life cycles. *Science*, 241: 1460 – 1466.
- Rudnik, D.T., R. Elmgren and J.B. Frithsen, 1985. Meiofaunal prominence and benthic seasonality in a coastal marine ecosystem. *Oecologia*, 67: 157 – 168.
- Russell, B.C., 1977. Population and standing crop estimates for rocky reef fishes of north-eastern New Zealand. *N.Z. Journal of Marine and Freshwater Research*,

11(1): 23 – 36.

- Saggiomo, V., A. Goffart, G.C. Carrada and J.H. Hecq, 1994. Spatial patterns of phytoplanktonic pigments and primary production in a semi-enclosed periantarctic ecosystem: the Strait of Magellan. *Journal of Marine Systems*, 5: 119 – 142.
- Sainsbury, K.J., 1987. Assessment and management of the demersal fishery on the continental shelf of Northwestern Australia. In: Polovina, J.J., and S. Ralston (ed.s), *Tropical snappers and groupers: biology and fishery management*. Westview Press:London. pp: 465 – 503.
- Sainsbury, K., A.E. Punt and A.D.M. Smith, 2000. Design of operational management strategies for achieving fishery ecosystem objectives. *ICES Journal of Marine Science*, 57: 731 – 741.
- Sakshaug, E., A. Bjorge, B. Gulliksen, H. Loeng and F. Mehlum, 1994. Structure, biomass distribution and energetics of the pelagic ecosystem in the Barents Sea – a synopsis. *Polar Biology*, 14(6): 405 – 411.
- Sala, E., 1997. Fish predators and scavengers of the sea urchin *Paracentrotus lividus* in protected areas of the north-west Mediterranean Sea. *Marine Biology*, 129: 531 – 539.
- Samoilys, M.A. and L.C. Squire, 1994. Preliminary observations on the spawning behaviour of coral trout, *Plectropomus leopardus* (Pisces: Serranidae), on the Great Barrier Reef. *Bulletin of Marine Science*, 54: 332 - 342.
- Santos, P.J.P., J. Castel and L.P. Souza-Santos, 1996. Seasonal variability of meiofaunal abundance in the oligo-mesohaline area of the Gironde estuary, France. *Estuarine, Coastal and Shelf Science*, 43: 549 – 563.
- Santschi, P.H., 1995. Seasonality in nutrient concentrations in Galveston Bay. *Marine*

- Environmental Research, 40: 337 – 362.
- Sarnelle, O., 1994. Inferring process from pattern: trophic level abundances and imbedded interactions. *Ecology*, 75: 1835 – 1841.
- Schmid, T. H., F.L. Murru and F. McDonald, 1993. Feeding habits and growth rates of bull (*Carcharinas leucas*), sandbar (*Carcharinas plumbeus*), sandtiger (*Eugomphodus taurus*), and nurse (*Ginglymostoma cirratum*) sharks. *Journal of Aquaculture and Aquatic Sciences*, 5(4): 100-105.
- Schreiber, R.A. and J.R. Pennock, 1995. The relative contribution of benthic microalgae to total microalgal production in a shallow sub-tidal estuarine environment. *Ophelia*, 42: 335 – 352.
- Schwinghamer, P., 1981. Characteristic size distribution of integral benthic communities. *Canadian Journal of Fisheries and Aquatic Sciences*, 38: 1255 – 1263.
- Sekine, M., H. Nakanishi, M. Ukita and S. Murakami (1991) A Shallow-Sea Ecological Model Using an Object-Oriented Programming Language. *Ecological Modelling*, 57: 221 – 236.
- Sharov, A.A., 1996. Modeling insect dynamics. In: E. Korpilahti, H. Mikkela, and T. Salonen (eds.) *Caring for the forest: research in a changing world*. Congress Report, Vol. II., IUFRO XX World Congress, 6-12 August 1995, Tampere, Finland. Gummerus Printing, Jyväskylä (Finland). pp. 293 – 303.
- Sheldon, R.W., A. Prakash and W.H. Sutcliffe Jr., 1972. The size distribution of particles in the ocean. *Limnology and Oceanography*, 17: 327 – 340.
- Shepherd, S.A. and I.M. Thomas, 1982. *Marine invertebrates of southern Australia*. D.J. Woolman, Government Printer, South Australia
- Siljak, D.D., 1974. Connective stability of complex ecosystems. *Nature*, 249: 280.

- Siljak, D.D., 1976. When is a complex ecosystem stable? *Mathematical Biosciences*, 25: 25 – 50.
- Silvert, W.L., 1981. Principles of ecosystem modelling. In: A.R. Longhurst (ed), *Analysis of marine ecosystems*. Academic Press: New York.
- Silvert, W. 1991. Experiments in benthic modelling. In: P. M. J. Herman and C. H. R. Heip (ed.s), *Report of the workshop Modelling the Benthos, Yerseke, Netherlands, 20-22 March 1991*. Delta Institute for Hydrobiological Research, Royal Netherlands Academy of Arts and Sciences, Comm. No. 538, pp. 175 – 188.
- Silvert, W.L., 1996a. Complexity. *Journal of Biological Systems*, 4: 585 – 591.
- Silvert, W.L., 1996b. Size-aggregation in models of aquatic ecosystems. *The Science and the Total Environment*, 183: 107 – 114.
- Silvertsen, K., 1997. Geographic and environmental factors affecting the distribution of kelp beds and barren grounds and changes in biota associated with kelp reduction at sites along the Norwegian coast. *Canadian Journal of Fisheries and Aquatic Sciences*, 54: 2872 – 2887.
- Skellam, J.G., 1951. Random dispersal in theoretical populations. *Biometrika*, 38: 196 – 218.
- Skovgaard, A., 1996. Mixotrophy in *Fragilidium subglobosum* (Dinophyceae) – growth and grazing responses as functions of light intensity. *Marine Ecology Progress Series*, 143: 247 – 253.
- Smayda, T.J., 1983. The phytoplankton of estuaries. In: B.H. Ketchum (ed.), *Estuaries and enclosed seas*. Elsevier:Amsterdam. P: 65 – 102.
- Smayda, T.J., 1997. Harmful algal blooms – their ecophysiology and general relevance to phytoplankton blooms in the sea. *Limnology and Oceanography*,

45: 1137 – 1153.

Smith, D.C. and C.M. MacDonald (eds.), 1997. King George whiting – 1996.

Compiled by the Bay & Inlet Fisheries and Stock Assessment Group. Fisheries

Victoria Assessment Report No 15. Fisheries Victoria: East Melbourne.

Smith, A.D.M., K.J. Sainsbury and R.A. Stevens (1999) Implementing effective

fisheries management systems – management strategy evaluation and the

Australian partnership approach. ICES Journal of Marine Science, 56: 697 –

979.

Soetaert, K. and P.M.J. Herman, 1994. One foot in the grave: zooplankton drift into

the Westerschelde estuary (the Netherlands). Marine Ecology Progress Series,

105: 19 – 29.

Soetaert, K., M. Vincx, J. Wittoeck, M. Tulkens and D. Van Gansbeke, 1994. Spatial

patterns of Westerschelde meiobenthos. Estuarine, Coastal and Shelf Science,

39: 367 – 388.

Sparre, P., 1991. An introduction to multispecies virtual population analysis. ICES

Marine Science Symposia, 193: 12 – 21.

Steele, J.H., 1998. From carbon flux to regime shift. Fisheries Oceanography, 7: 176 –

181.

Steele, J.H. and I.E. Baird, 1968. Production ecology of a sandy beach. Limnology

and Oceanography, 13: 14 – 25.

Steele, J.H. and E.W. Henderson, 1992. The role of predation in plankton models.

Journal of Plankton Research, 14: 157 – 172.

Steele, J.H. and E.W. Henderson, 1995. Predation control of plankton demography.

ICES Journal of Marine Science, 52: 565 – 573.

Steimle, F.W. Jr., 1982. The benthic macroinvertebrates of the Block Island Sound.

- Estuarine, Coastal and Shelf Science, 15: 1 – 16.
- Steimle, F.W. Jr., 1985. Biomass and estimated productivity of the benthic macrofauna in the New York Bight: A stressed coastal area. *Estuarine, Coastal and Shelf Science*, 21: 539 – 554.
- Sterner, R.W., J.J. Elser, T.H. Chrzanowski, J.H. Schampel and N.B. George, 1996. Biogeochemistry and trophic ecology: a new food web diagram. In: G.A. Polis and K.O. Winemiller (ed.s), *Food webs: integration of patterns and dynamics*. Chapman and Hall, New York, London.
- Stickney, H.L., R.R. Hood and D.K. Stoecker, 2000. The impact of mixotrophy on planktonic marine ecosystems. *Ecological Modelling*, 125: 203 – 230.
- Stoecker, D.K., 1999. Mixotrophy among dinoflagellates. *Journal of Eukaryotic Microbiology*, 46: 397 – 401.
- Sugihara G., S. Garcia, J.A. Gulland, J.H. Lawton, H. Maske, R.T. Paine, T. Platt, E. Rachor, B.J. Rothschild, E.A. Ursin and B.K.K. Zeitzschel, 1984. *Ecosystem Dynamics: Group Report*. In: R.M. May (ed), *Exploitation of Marine Communities*, Dahlem Konferenzen 1984. Springer-Verlag:Berlin, Heidelberg, New York, Tokyo. pp 131 – 153.
- Talley, T.S., P.K. Dayton and S.E. Ibarra-Obando, 2000. Tidal flat macrofaunal communities and their associated environments in estuaries of southern California and northern Baja California, Mexico. *Estuaries*, 23: 97 – 114.
- Taylor, C.D. and B.L. Howes, 1994. Effect of sampling frequency on measurements of seasonal primary production and oxygen status in near-shore coastal ecosystems. *Marine Ecology Progress Series*, 108: 193 – 203.
- Tett, P and H. Wilson, 2000. From biogeochemical to ecological models of marine microplankton. *Journal of Marine Systems*, 25: 431 – 446.

- Thayer, G.W., D.R. Colby and W.F. Hettler Jr., 1987. Utilization of the red mangrove prop root habitat by fishes in south Florida. *Marine Ecology Progress Series*, 35: 25 – 38.
- Thomas, H., V. Ittekkot, C. Osterroht and B. Schneider, 1999. Preferential recycling of nutrients – the ocean's way to increase new production and to pass nutrient limitation? *Limnology and Oceanography*, 44: 2005 – 2011.
- Tilman, D. and J.A. Downing, 1994. Biodiversity and stability in grasslands. *Nature*, 367: 363 – 365.
- Tober, J., C. Fritz, E. LaBrecque, P.J. Behr and I. Valiela, 1996. Abundance, biomass, species richness of fish communities in relation to Nitrogen-loading rates of Waquiot Bay estuaries. *Biological Bulletin*, 191: 321 – 322.
- Tremblay, J.É., L. Legendre and J.C. Therriault, 1997. Size-differential effects of vertical stability on the biomass and production of phytoplankton in a large estuarine system. *Estuarine, Coastal and Shelf Science*, 45: 415 – 431.
- Turner, J.T., S.F. Bruno, R.J. Larson, R.D. Staker and G.M. Sharma, 1983. Seasonality of plankton assemblages in a temperate estuary. *P.S.Z.N. Marine Ecology*, 4: 81 – 99.
- Ulanowicz, R.E., 1986. *Growth and development: ecosystem phenomenology*. Springer-Verlag: New York.
- Valentine, J.F. and K.L. Heck, 1993. Mussels in seagrass meadows: their influence on macroinvertebrate abundance and secondary production in the northern Gulf of Mexico. *Marine Ecology Progress Series*, 96: 63 – 74.
- Valiela, I., K. Foreman, M. LaMontagne, D. Hersh, J. Costa, P. Peckol, B. DeMeo-Andreson, C. D'Avanzo, M. Babione, C. Sham, J. Brawley and K. Lajtha, 1992. *Couplings of watersheds and coastal waters: sources and consequences of*

- nutrient enrichment in Waquoit Bay, Massachusetts. *Estuaries*, 15: 443 – 457.
- Valiela, I., J. McClelland, J. Hauxwell, P.J. Behr, D. Hersh and K. Foreman, 1995. Macroalgal blooms in shallow estuaries: Controls and ecophysiological and ecosystem consequences. *Limnology and Oceanography*, 42: 1105 – 1118.
- Varela, R.A., A. Cruzado and J.E. Gabaldón, 1995. Modelling primary production in the North Sea using the European Regional Seas Ecosystem Model. *Netherlands Journal of Sea Research*, 33: 337 – 361.
- van Tongeren, O.F.R., 1995. Data analysis or simulation model: a critical evaluation of some methods. *Ecological Modelling*, 78: 51 – 60.
- Vanhove, S., H.J. Lee, M. Beghyn, D. Van Gansbeke, S. Brockington and M. Vincx, 1998. The metazoan meiofauna in its biogeochemical environment: the case of an Antarctic coastal sediment. *Journal of the Marine Biology Association of the United Kingdom*, 78: 411 – 434.
- Ver, L.M.B., F.T. Mackenzie and A. Lerman, 1999. Biogeochemical responses of the carbon cycle to natural and human perturbations: Past, present, and future. *American Journal of Science*. 299(7-9): 762 – 801.
- Waide, J.B. and J.R. Webster, 1976. Engineering systems analysis: applicability to ecosystems. In: B.C. Patten (ed), *Systems Analysis and Simulation in Ecology*. Academic Press: New York.
- Walker, S.J., 1997. Hyrdodynamic models of Port Phillip Bay. Technical Report No. 38. Port Phillip Bay Environmental Study. CSIRO: Canberra, Australia.
- Walker, S.J., 1999. Coupled hydrodynamic and transport models of Port Phillip Bay, a semi-enclosed bay in south-eastern Australia. *Marine and Freshwater Research*, 50: 469 – 482.
- Walker, D.I. and A.J. McComb, 1992. Seagrass degradation in Australian coastal

- waters. *Marine Pollution Bulletin*, 25: 191 - 195.
- Walker, S.J. and C.R. Sherwood, 1997. A Transport Model of Port Phillip Bay. Technical Report No. 39. Port Phillip Bay Environmental Study. CSIRO: Canberra, Australia.
- Walters, C., 1998. ECOSIM and ECOSPACE: basic considerations. In: D. Pauly (ed), Use of ECOPATH and ECOSIM to Evaluate Strategies for Sustainable Exploitation of Multi-species Resources: Proceedings of a Workshop held March 25 – 27, 1998. Fisheries Centre, University of British Columbia: Vancouver, Canada.
- Walters, C., V. Christensen and D. Pauly, 1997. Structuring dynamic models of exploited ecosystems from trophic mass-balance assessments. *Reviews in Fish Biology and Fisheries*, 7: 139 – 172.
- Walters, C., D. Pauly and V. Christensen, 1999. ECOSPACE: prediction of mesoscale spatial patterns in trophic relationships of exploited ecosystems, with emphasis on the impacts of marine protected areas. *Ecosystems*, 2(6): 539 – 554.
- Walters, C., D. Pauly, V. Christensen and J.F. Kitchell, 2000. Representing density dependent consequences of life history strategies in an ecosystem model: ECOSIM II. *Ecosystems*, 3: 70 – 83.
- Wantiez, L., M. Harmelin-Vivien and M. Kulbicki, 1996. Spatial and temporal variation in a soft-bottom fish assemblage in St. Vincent Bay, New Caledonia. *Marine Biology*, 125: 801 – 812.
- Wantiez, L., 1996. Comparison of fish assemblages sampled by a shrimp trawl and a fish trawl in St Vincent Bay, New Caledonia. *Journal of the Marine Biology Association of the United Kingdom*, 76: 759 – 775.
- Wantiez, L., 1998. Structure des communautés de poissons des fonds meubles du lagon

- nord de nouvelle-calédonie et leurs relations avec le milieu. *Cybiu*, 22(2): 107 – 122.
- Warburton, K., 1979. Growth and production of some important species of fish in a Mexican coastal lagoon system. *Journal of Fish Biology*, 14: 449 – 464.
- Warwick, R.M. and R. Price, 1975. Macrofauna production in an estuarine mud-flat. *Journal of the Marine Biology Association of the United Kingdom*, 55: 1 – 18.
- Watt, K.E.F., 1975. Critique and comparison of biome ecosystem modeling. In: B.C. Patten (ed), *System analysis and simulation in ecology*, III. Academic Press: New York, London. pp: 139 – 152.
- Webber, H.H. and H.V. Thurman, 1991. *Marine Biology*; 2nd ed. Harper Collins Publishers: New York
- Weins, J.A., 1989. Spatial scaling and ecology. *Functional Ecology*, 3: 385 – 397.
- Welsh, B.L., 1980. Comparative nutrient dynamics of marsh-mudflat ecosystem. *Estuarine, Coastal and Shelf Science*, 10: 143 – 164.
- Whitfield, A.K., 1993. Fish biomass estimates from the littoral zone of an estuarine coastal lake. *Estuaries*, 16: 280 – 289.
- Wiegert, R.G., 1975. Simulation modeling of the algal-fly components of a thermal ecosystem: effects of spatial heterogeneity, time delays, and model condensation. In: B.C. Patten (ed), *System analysis and simulation in ecology*, III. Academic Press: New York, London. pp: 157 – 181.
- Wiegert, R.G., 1977. A model of a thermal spring food chain. In: C. Hall and J.W. Day (ed.s), *Ecosystem Modelling In Theory and Practice*. Wiley and Sons: New York, London, Sydney. pp: 290 – 315.
- Williams, D.McB. and A.I. Hatcher, 1983. Structure of fish communities on outer slopes of inshore, mid-shelf and outer-shelf reefs of the Great Barrier Reef.

- Marine Ecology Progress Series, 10: 239 – 250.
- Wilson, R.S., B. F. Cohen and G.C.B. Poore, 1993. The Role of Suspension-feeding and Deposit-feeding Benthic Macroinvertebrates in Nutrient Cycling in Port Phillip Bay. Technical Report No. 10. Port Phillip Bay Environmental Study. CSIRO: Canberra, Australia.
- Wilson, R.S., S. Heislors and G.C.B. Poore, 1998. Changes in benthic communities of Port Phillip Bay, Australia, between 1969 and 1995. *Marine and Freshwater Research*, 49: 847 – 861.
- Wright, B.A. and L. Hulbert, 2000. Shark abundance increases in the Gulf of Alaska. *PICES Press*, 8(2)
- Wolff, M., 1994. A trophic model for Tongoy Bay – a system exposed to suspended scallop culture (Northern Chile). *Ecological Modelling*, 182: 149 – 168.
- Yachi, S. and M. Loreau, 1999. Biodiversity and ecosystem functioning in fluctuating environments: the insurance hypothesis. *Proceedings of National Academy of Sciences USA* 96: 1463 – 1468.
- Yearsley, J.R. and Lettenmaier D.P., 1987. Model complexity and data worth: An assessment of changes in the global carbon budget. *Ecological Modelling*, 39: 201-226.
- Yodzis, P., 1981. The stability of real ecosystems. *Nature*, 289: 674 – 676.
- Yodzis, P., 2000. Diffuse effects in food webs. *Ecology*, 81: 261 – 266.
- Yodzis, P., 2001a. Must top predators be culled for the sake of fisheries? *Trends in Ecology and Evolution*, 16: 78 – 84.
- Yodzis, P., 2001b. Culling predators to protect fisheries: a case of accumulating uncertainties – response. *Trends in Ecology and Evolution*, 16: 282 – 283.
- Yool, A., 1998. The Dynamics of Open-Ocean Plankton Ecosystem Models. PhD

Thesis, Ecology and Epidemiology, Department of Biological Sciences,
University of Warwick. pp 354.

Young, P., S. Parkinson and M. Lees, 1996. Simplicity out of complexity in
environmental modelling: Occam's razor revisited. *Journal of Applied
Statistics*, 23: 165 – 210.

Zeigler, B.P., 1976. The aggregation problem. In: B.C. Patten (ed), *System analysis
and simulation in ecology*, IV. Academic Press: New York, London. pp: 299 –
311.